



## **Towards CHAOS-5 - How can Swarm contribute?**

**Finlay, Chris; Olsen, Nils; Tøffner-Clausen, Lars**

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Finlay, C., Olsen, N., & Tøffner-Clausen, L. (2014). Towards CHAOS-5 - How can Swarm contribute? *Geophysical Research Abstracts*, 16, [EGU2014-10878]. <http://adsabs.harvard.edu/abs/2014EGUGA..1610878F>

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## **Towards CHAOS-5 - How can Swarm contribute?**

Christopher C. Finlay, Nils Olsen, and Lars Tøffner-Clausen

Division of Geomagnetism, DTU Space, Technical University of Denmark (cfinlay@space.dtu.dk)

The launch of ESA's satellite trio Swarm in November 2013 opens an exciting new chapter in the observation and monitoring of Earth's magnetic field from space. We report preliminary results from an extension of the CHAOS series of geomagnetic field models to include both scalar and vector field observations from the three Swarm satellites, along with the most recent quasi-definitive ground observatory data. The fit of this new update CHAOS field model to the Swarm observations will be presented in detail providing useful insight the initial Swarm data. Enhancements of the CHAOS modelling scheme include a 1 minute time resolution for the RC index and anisotropic weighting of vector field data depending on quasi-dipole latitude. We shall also report on the perspective given by the initial Swarm data on rapid field changes currently taking place in the Atlantic sector.